

HEPATITIS A

General Information

What is hepatitis?

“Hepatitis” means inflammation of the liver. The liver is a vital organ that processes nutrients, filters the blood, and fights infections. When the liver is inflamed or damaged, its function can be affected.

Hepatitis is most often caused by a virus. In the United States, the most common types of viral hepatitis are Hepatitis A, Hepatitis B, and Hepatitis C. Heavy alcohol use, toxins, some medications, and certain medical conditions can also cause hepatitis.

What is Hepatitis A?

Hepatitis A is a highly contagious liver infection caused by the Hepatitis A virus. It can range in severity from a mild illness lasting a few weeks to a severe illness lasting several months.

Who is at risk?

Although anyone can get Hepatitis A, some people are at greater risk, such as those who:

- Travel to or live in countries where Hepatitis A is common
- Use recreational drugs, whether injected or not
- Have sexual contact with someone who has Hepatitis A
- Have clotting-factor disorders, such as hemophilia
- Are men who have sexual encounters with other men
- Are household members or caregivers of a person infected with Hepatitis A

How common is Hepatitis A?

Hepatitis A still occurs in the United States, although not as frequently as it once did. Over the last several decades, there has been more than a 90% decrease in Hepatitis A cases. New cases are now estimated to be around 3,000 each year. Many experts believe this decline is a result of the vaccination of children and people at risk for Hepatitis A. Many of the new cases, however, are from American travelers who got infected while traveling to parts of the world where Hepatitis A is common.



Hepatitis A can be prevented with a safe and effective vaccine.

How is Hepatitis A spread?

Hepatitis A is usually spread when a person ingests fecal matter—even in microscopic amounts—from contact with objects, food, or drinks contaminated by feces or stool from an infected person.

Hepatitis A can be spread when:

- An infected person does not wash his/her hands properly after going to the bathroom and then touches objects or food
- A caregiver does not properly wash his or her hands after changing diapers or cleaning up the stool of an infected person
- Someone engages in sexual activities with an infected person

Hepatitis A also can be spread through contaminated food or water. Contamination of food can happen at any point: growing, harvesting, processing, handling, and even after cooking. This most often occurs in countries where Hepatitis A is common.

Continued on next page



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

What are the symptoms of Hepatitis A?

Not everyone has symptoms. If symptoms develop, they usually appear 2 to 6 weeks after infection and can include:

- Fever
- Vomiting
- Grey-colored stools
- Fatigue
- Abdominal pain
- Joint pain
- Loss of appetite
- Dark urine
- Jaundice
- Nausea

Symptoms are more likely to occur in adults than in children. They usually last less than 2 months, although some people can be ill for as long as 6 months.



People can spread Hepatitis A even if they don't look or feel sick. Many children and some adults have no symptoms.

How is Hepatitis A diagnosed and treated?

A doctor can determine if a person has Hepatitis A by discussing his or her symptoms and taking a blood sample. To treat Hepatitis A, doctors usually recommend rest, adequate nutrition, fluids, and medical monitoring. Some people will need to be hospitalized. It can take a few months before people begin to feel better.

How serious is Hepatitis A?

Most people who get Hepatitis A feel sick for several months, but they usually recover completely and do not have lasting liver damage. Sometimes Hepatitis A can cause liver failure and death, although this is rare and occurs more commonly in people older than 50 and people with other liver diseases.

Can Hepatitis A be prevented?

Yes. The best way to prevent Hepatitis A is by getting vaccinated. Experts recommend the vaccine for all children, and people with certain risk factors and medical conditions. The vaccine is also recommended for travelers to certain international countries, even if travel occurs for short times or on closed resorts. The Hepatitis A vaccine is safe and effective and given as 2 shots, 6 months apart. Both shots are needed for long-term protection. Ask if your health plan will cover travel related vaccines. You can get vaccinated at your doctor's office, as well as travel clinics and other locations. Lower cost vaccination may be available at certain pharmacies and your local health department.

Who should get vaccinated against Hepatitis A?

Vaccination is recommended for certain groups, including:

- All children at age 1 year
- Travelers to countries where Hepatitis A is common
- Family and caregivers of adoptees from countries where Hepatitis A is common
- Men who have sexual encounters with other men
- Users of recreational drugs, whether injected or not
- People with chronic or long-term liver disease, including Hepatitis B or Hepatitis C
- People with clotting-factor disorders

For more information

Talk to your health professional, call your health department, or visit www.cdc.gov/hepatitis or www.cdc.gov/travel.

Help Prevent the Spread of Hepatitis A

1: CLEAN UP SURFACES



2: DISINFECT SURFACES



3: WASH YOUR HANDS



Scientific experts from the U.S. Centers for Disease Control and Prevention (CDC) helped to develop this poster. For more information on Hepatitis A prevention, please see <https://www.cdc.gov/hepatitis/hav/>



Hepatitis A is going around.

Hepatitis A is a virus that can cause liver disease.

Protect yourself: Get your Hepatitis A vaccination

Who should get vaccinated?

- People currently experiencing homelessness
- People who use street drugs
- Men who have sex with men
- People recently in jail or prison

You can also protect yourself and others by washing your hands ANYTIME you:

- Go to the bathroom
- Touch, eat, or serve food or drink
- Use street drugs of any kind

